

IRON CONTAMINATION DURING COMMERCIAL GRINDING OF SPICES

JANITHA P. PANDUWAWALA, CHAMARA D.K. ILLEPERUMA AND
U. SAMARAJEEWA

*Department of Food Science & Technology, University of Peradeniya,
Peradeniya, Sri Lanka.*

(Date of receipt : 29 May 1987)

(Date of acceptance : 06 January 1988)

Abstract : The concentration of copper, iron and zinc were estimated by atomic absorption spectrophotometry in powdered spices purchased from retailers. The three elements were compared by grinding the whole spices by mortar and pestle, grinding stone, food blender and in commercial mills. The iron, due to wear and tear of the machinery, contaminate the spices during commercial grinding increasing the iron content 3-5 fold; the concentrations being as high as 750 $\mu\text{g/g}$ of ground spices. Highest contamination were seen in turmeric powder. The iron contaminated spices are ferromagnetic confirming the particulate nature of iron in spices. The copper and zinc content were comparable in the replicates ground by different methods as well as in the samples purchased from retailers. Further study on the nutritional significance of iron contamination in ground spices appears warranted.

1. Introduction

Spices are added to enhance the flavour, aroma and colour of food but are not considered as sources of minerals or other nutrients. Ground spices are reported to contain high iron concentrations, sometimes 3-4 times higher than in other edible plant components, implicating possible nutritional benefits through the spices.^{3,6} However, the origin of such high levels of iron is not understood.

In this study, experiments were directed at determining the origin of the copper, iron and zinc content of ground spices.

2. Materials and Methods

2.1 Spices

The following whole or commercially ground spices and curry powder were purchased from the retailers in Gampaha, Kandy, Matale, Nugegoda, Peradeniya and Warakapola.

- Pods of ripe chillie (*Capsicum annum*)
- Berries of pepper (*Piper nigrum*)
- Rhizomes of turmeric (*Curcuma longa*)
- Seeds of coriander (*Coriandrum sativum*)
- Fruits of cumin (*Cuminum cyminum*)

2.2 Grinding

Whole spices were washed with demineralized water, and dried at 70°C in the vacuum oven. Replicate samples of the dried spices were ground by the following methods:

- (a) In a laboratory mortar and pestle
- (b) In a clean grinding stone at kitchen
- (c) In a food blender (National Model MX-11PN)
- (d) In commercial mills in Kandy

Estimations were made from triplicate samples ground by each method.

2.3 Estimation of copper, iron and zinc

The powdered spices (1g) were ignited in a muffle furnace at 460°C for eight hours. This was moistened with glass distilled demineralized water (2 ml) and concentrated nitric acid (5 ml) and warmed for few minutes over the flame. The solutions were filtered in to 100 ml volumetric flasks through Whatman No. 541 filter paper and made up to the mark.

The estimations were carried out using a Perkin-Elmer Model 2380 Atomic Absorption Spectrophotometer containing an adjustable titanium burner head and air/acetylene flame, using high intensity "Intensitron" lamps at the wave lengths and slit widths given in Table 1.⁷

Table 1

element	wave length (nm)	slit (nm)
copper	324.8	0.7
iron	248.3	0.2
zinc	213.9	0.7

The quantitative analysis of the elements were conducted using standard curves obtained using the same instrument, for standard solutions containing the respective cations.² The coefficient of variation for the determinations varied up to 15% for the home ground samples and up to 40% for

the mill ground samples; the values for commercially purchased samples varied between 15–79%.

2.4 Ferromagnetism

The powdered spices were firmly packed to a length of 4 cm in a glass tube of 10 cm x 10 mm (internal diameter) and suspended in a double pan laboratory balance at the centre of an electromagnet set underneath the balance. The tube was balanced using the weights.

A magnetic field strength of up to 3000 oersted was produced by the electromagnet with pole gap of 2 cm, resistance of 15.3 ohm and 2337 turns which was supplied with a power of 22 volts and 4.5 amp to produce a magnetic flux of 60 maxwell turns using Eleminac type B 2027 generator. The magnetic flux produced was measured by a calibrated fluxmeter in maxwell turns per 100 cm² search coil.⁵

3. Results and Discussion

3.1 Commercially ground spices

The powdered spices purchased from the retailers contained high iron content as already reported⁶ and the copper and zinc contents were much less than iron (Table 2). The iron content varied widely among the samples with standard deviations up to 70% of the mean whereas the standard deviations for copper and zinc contents were less than 30% of the mean. The high inhomogeneity of iron content in commercial spice powders suggest the presence of iron as a contaminant rather than a naturally occurring constituent of the raw materials.

The iron content in the commercial powders did not show any correlation with the location of purchase. Of the different types of spices examined turmeric had the highest iron content (Table 2).

3.2 Home ground spices

The spices ground by three methods at home/laboratory showed comparable figures for copper, iron and zinc content (Table 3). The iron content found in the home ground spices was only 25 to 35 per cent of the levels reported in commercially ground spices. In contrast to the iron content, the copper and zinc contents were the same in both home ground and commercially purchased spices further indicating the high iron content to be a contamination during grinding in commercial mills.

Table 2. Copper, iron and zinc concentrations ($\mu\text{g/g}$) in commercial powdered spices purchased from retailers.

Spice element	No. of samples	Mean	S.D.	Range
Chillie	25			
copper		36.3	14.4	22-64
iron		458.0	144.2	214-806
zinc		20.0	7.3	10-30
Curry	16			
copper		33.1	5.8	22-41
iron		608.1	396.1	197-1998
zinc		29.0	4.4	24-40
Pepper	5			
copper		52.8	11.8	38-71
iron		449.4	326.0	100-912
zinc		19.4	8.4	8-28
Turmeric	15			
copper		17.3	13.7	1-47
iron		740.6	491.4	107-1617
zinc		27.1	12.7	6-60

Table 3. Copper, iron and zinc concentrations ($\mu\text{g/g}$) in spices ground using mortar and pestle, grinding stone or food blender.

Spice	element	Method of grinding			Mean \pm S.D.
		mortar & pestle	grinding stone	food blender	
Chillie (3)					
	copper	49	43	41	44.3 \pm 3.3
	iron	126	138	133	132.3 \pm 4.9
	zinc	18	16	12	15.3 \pm 2.4
Coriander (3)					
	copper	41	38	35	38.3 \pm 2.4
	iron	143	146	147	145.3 \pm 1.6
	zinc	32	30	28	30.0 \pm 1.6
Cumin (3)					
	copper	49	49	44	47.3 \pm 2.3
	iron	170	178	167	171.6 \pm 4.6
	zinc	30	28	25	27.6 \pm 2.0
Pepper (3)					
	copper	54	52	54	53.3 \pm 0.9
	iron	74	70	68	70.6 \pm 2.4
	zinc	24	25	28	25.6 \pm 1.6
Turmeric (3)					
	copper	30	25	24	26.3 \pm 2.6
	iron	130	143	140	137.6 \pm 5.5
	zinc	28	28	29	28.3 \pm 0.4

Results for the same samples ground in the mill.

Among the spices cumin seed is reported to be a naturally rich source of iron⁴ containing 950 $\mu\text{g/g}$. The samples examined in this study contained only 170 $\mu\text{g/g}$ of iron in cumin seeds.

3.3 Commercial grinders

When replicates from the whole spices ground at home were ground in commercial grinders, a 3–5 fold increase in the iron content was noted whereas the copper and zinc contents were the same as in home ground spices (Figure 1) confirming that iron contamination occurs during grinding. The added iron content varied from mill to mill widely. Of the 10 grinding mills compared for added iron content (Table 4) the mill No. 1 exhibited highest contamination for all types of spices.

The grinders used in Sri Lanka are of “burr type” and consist of two roughened cast iron discs, where flutes are grooved in the two facing sides. One plate rotates on a shaft and the whole spices are fed between the plates for grinding by crushing and shearing action. It is quite possible that iron, in a fine particulate form, is produced due to wear and tear during grinding. The higher iron content noted with turmeric (Table 2 and 4) is probably due to higher hardness of turmeric rhizomes causing greater wear and tear compared with the other spices. It is also possible that higher iron contamination occur in grinding mills where the plates are new and rough.

3.4 Magnetic properties

If iron contamination occurs in spices during grinding, the particulate nature of iron should add a ferromagnetic effect to the ground spices when tested by Gouy method.⁵ Home ground spices when subjected to a magnetic field were diamagnetic, exerting no force whereas the commercially ground spices exhibited a positive force on the balance confirming the particulate nature of iron. However, the ferromagnetic force was not sufficiently strong to be measured under our experimental conditions.

3.5 Nutritional significance

In cooking foods the total amount of spices added average to 5–10 g per person with a possible daily contribution of 5 mg iron per person from spices containing about 500 $\mu\text{g/g}$ of iron, as observed in this study. This falls very close to the recommended daily dietary allowance of 6–15 mg of iron.⁸ However, iron existing in the ferric state in the powdered spices may not be readily available for absorption during digestion.¹⁰ The bioavailability of iron depends on several parameters. On one hand the presence of ascorbic acid and citric acid contributed from lime during cooking of vegetables may convert the iron into chelate forms that could be readily absorbed. On the




Figure 1. Copper, iron and zinc concentrations ($\mu\text{g/g}$) in spices ground at home and commercial mills from the same samples of whole spices. Figures on bars refer to the number of samples assayed.

Table 4. Iron, Copper and Zinc concentrations ($\mu\text{g/g}$) in the same chillie, coriander, cumin and turmeric samples ground in different commercial grinders. They are replicates of samples ground by home methods (Table 2).

Mill No.	Chillie		Coriander		Cumin		Turmeric					
	Iron	Copper Zinc	Iron	Copper Zinc	Iron	Copper Zinc	Iron	Copper Zinc				
	1	743	34 16	866	49 26	932	25 20	1079	34 28			
2	387	44 16	270	45 24	483	34 19	279	36 24				
3	632	44 14	373	45 24	538	48 21	779	26 27				
4	244	46 13	554	35 24	308	41 23	696	26 27				
5	641	45 14	474	49 26	329	39 22	768	30 25				
6	279	44 18										
7	275	43 13										
8	444	40 14										
9	371	46 11										
10	267	45 13										
Mean	429	43.1 14.6	507	44.6 24.8	518	37.4 21	720	30.4 26.2				
S.D.	182	3.6 1.9	227	5.7 1.0	251	8.5 1.5	287.1	4.5 1.6				
Range	244-743	34-46	13-18	270-866	35-49	24-26	308-932	25-48	19-23	279-1079	26-36	24-28

other hand the iron may be complexed by phytates or oxalates or made insoluble by phosphates making them non-available for absorption. Consumption of contaminated foods that provide 100 mg iron per day is reported to cause siderosis in the Bantu populations.¹⁰ The spices cannot contribute such high iron levels to foods.

Iron in the ferric state may also produce negative effects on foods by oxidizing the anthocyanin pigments, which contribute colour to the foods and oxidizing ascorbic acid and other antioxidants in foods. The possible implications of the iron contamination from grinding mills on the nutrition need be examined further as spices are consumed considerably by Sri Lankans.

Zinc at concentrations above 40 mg/kg in water is reported to cause adverse physiological reactions in humans.⁹ The maximum acceptable daily intake of copper is 0.5 mg/kg.⁹ The contribution of zinc and copper to foods by the spices is much lower and could not be hazardous.

4. Conclusions

Contamination of iron due to wear and tear of the parts of mills cause 3–5 fold increase in iron content, in commercially ground spices. The iron appears in the particulate form in the spice powders. Further study on the nutritional significance of iron contamination in ground spices appears warranted.

Acknowledgements

We wish to thank Dr. M.A. Careem of Department of Physics, University of Peradeniya, for providing the facilities for magnetic measurements and Mrs. C. Baranage of the Food Testing Laboratory, Municipal Council, Kandy for providing commercially ground spices.

REFERENCES

1. ANONYMOUS (1970). *Fourteenth report of the joint FAO/WHO Expert Committee on Food Additives*. Geneva 36 p.
2. CHAPMAN, H.H. & PRATT, P.F. (1961) *Methods of analysis of soils, plants and water*. University of California, Division of Agricultural Sciences. p. 161–175.
3. CHRISTENSEN, R.E., BECKMAN, R.M. & BIRDSALL, J.J. (1968) *J. Assoc. Offic. Anal. Chem.*, 51: 1003.

4. FOX, SUSAN, J. & BENDER, A.E. (1977) *J. Food Technol.*, **12**: 535–539.
5. LEWIS, J. & WILKINS, R.G. (1960) *Modern Co-ordination Chemistry* p. 400, Wiley Interscience Publishers, New York.
6. PAUL, A.A. & SOUTHGATE, D.A.T. (1978) *The Composition of Foods*, Fourth revised and extended edition of M.R.C. Special Report No. 297: 271; Elsevier North-Holland Biomedical Press, Amsterdam.
7. PERKIN–ELMER (1976) *Analytical Methods for Atomic Absorption Spectrophotometry*, Norwalk, Connecticut, USA. p.1–12.
8. PYKE, M. (1982) *Success in Nutrition*, John Murray (Publishers) Ltd., London. 115–121.
9. STUTTS, V.J. (1981) Nutritional hazards, *In Food Safety*, edited by Robert, H.R. John Wiley & Sons, 113 p. New York, p. 113.
10. UNDERWOOD, E.J. (1977) *Trace Elements in Human and Animal Nutrition*, 25 p, Fourth Edition, Academic Press, London.