

RESEARCH NOTE

Effects of caffeine and a tea flavanoid on *Monacrosporium ambrosium*, ectosymbiote of shot-hole borer beetle of tea

Naleen B. Jayaratna, D. Nedra Karunaratne and N. Savitri Kumar*

Department of Chemistry, Faculty of Science, University of Peradeniya, Peradeniya.

Revised: 08 June 2008; Accepted: 20 July 2009

Keywords: Caffeine, ectosymbiote, flavanoid, *Monacrosporium ambrosium*, tea

Shot-hole borer (SHB) beetle *Euwallacea fornicatus* (*Xyleborus fornicatus*) Eichh. (Coleoptera: Scolytidae) is the most serious insect pest of tea (*Camellia sinensis* L.) plantations in Sri Lanka. Caffeine, the major alkaloid found in tea, inhibits the growth of both *Monacrosporium ambrosium*, ectosymbiote of SHB and the SHB beetle, in laboratory culture media^{1,2}. Caffeine contents of two tea cultivars that are resistant (TRI 2023) and susceptible (TRI 2025) to SHB infestation³, indicated that there was an increase in caffeine content after SHB infestation of both tea cultivars, the increase being higher in the more susceptible tea cultivar⁴. However, the two organisms are able to develop within tea stems that have a caffeine concentration which is higher than the concentration that inhibited the development of *M. ambrosium* and SHB beetles in laboratory culture media.

Flavan-3-ols and their gallate esters (catechins) may constitute up to 30% of young tea shoots. Epigallocatechin gallate (EGCG), the major phenolic compound found in tea, constitutes 9-13 % of the dry weight of young tea shoots⁵. The present study was carried out to determine whether EGCG modulates the effect of caffeine on *M. ambrosium* in laboratory culture media.

The effect of caffeine and EGCG on *M. ambrosium* was studied by determining the fungal colony diameter, sporulation and spore germination on solid agar media and mycelial mass in liquid culture media respectively, into which varying amounts of these two phytochemicals were incorporated. The ambrosia fungus was isolated¹ from a surface sterilized adult female SHB removed

from a gallery in a tea stem. The eight types of media (1-8), were prepared² in Cook's 2 quarter strength agar medium: 1. Control, 2. Caffeine 500 ppm, 3. EGCG 500 ppm, 4. Caffeine 500 ppm + EGCG 500 ppm, 5. Caffeine 1000 ppm, 6. Caffeine 1000 ppm + EGCG 500 ppm, 7. Caffeine 250 ppm and 8. Caffeine 250 ppm + EGCG 500 ppm. A fungal disc, cut from a well grown colony of *M. ambrosium* was placed in the center of a Petri dish containing each of the above media and the diameter of the fungal colony was measured daily for five days (3 replicates). Sporulation was studied using a spore suspension prepared from fungal discs (30, 8 mm diameter) cut from a fungal colony grown in each type of medium. These were mixed with sterile water (4 mL, Vibromix, 30 s), filtered through sterilized glass wool and the number of spores in each sample was counted using a hemocytometer (3 replicates). Spore germination was studied using fungal mycelia scraped from the tops of fungal colonies growing in different media. These were dissolved in sterile water (5 mL), shaken (Vibromix 30 s), filtered through glass wool and centrifuged (3 min, 3000 rpm). The supernatant (3 mL) was removed; one drop of solution from the bottom of the tube was placed on a microscope slide and left aside in a moisture chamber for 9 h. The number of germinated spores in 10 microscopic fields was counted. The percentage of spores that had germinated in each sample was calculated (3 replicates). Several experiments showed that sporulation did not occur in the presence of 1000 ppm of caffeine. Therefore, sporulation and spore germination in media 5 and 6 containing 1000 ppm of caffeine were not studied.

The effect of caffeine and EGCG on liquid cultures were studied by adding a spore suspension (20 mL,

*Corresponding author (nskumar@pdn.ac.lk)

5×10^3 spores/cm²)¹ to a Czapek-dox liquid culture medium (2000 mL)⁴. Calculated amounts of caffeine and/or EGCG was added to each of twenty 250 mL conical flasks under sterile conditions to obtain concentrations

similar to the solid agar media 1, 2, 3, 4 and the liquid culture medium (100 mL) was added to each flask and left on a shaker for 5 d. Fungal mycelia were separated and weighed (5 replicates).




Figure 1: Control (1), EGCG 500 ppm (3) and Caffeine 1000 ppm (5) Caffeine 1000 ppm + EGCG 500 ppm (6)



Figure 2: Control (1), Caffeine 500 ppm (2), EGCG 500 ppm (3) and caffeine 500 ppm + EGCG 500 ppm (4)

Table 1: Effect of Caffeine and EGCG on *M. ambrosium* on laboratory culture media

| Medium (No.) | Treatment Parameters | Growth Parameters- Percentage Reduction | | | |
|--------------|---------------------------------|---|-------------|-------------------|-------------|
| | | Fungal colony diameter | Sporulation | Spore germination | Fungal mass |
| 1 | Control | 0.0 | 0.0 | 0.0 | 0.0 |
| 2 | Caffeine 500 ppm | 17.5 | 85.0 | 55.0 | 23.6 |
| 3 | EGCG 500 ppm | 0.0 | 5.0 | 36.0 | 56.0 |
| 4 | Caffeine 500ppm + EGCG 500ppm | 15.0 | 90.0 | 39.0 | 58.0 |
| 5 | Caffeine 1000 ppm | 42.0 | Not done | Not done | Not done |
| 6 | Caffeine 1000ppm + EGCG 500 ppm | 37.0 | Not done | Not done | Not done |
| 7 | Caffeine 250 ppm | 7.5 | 70 | 55.0 | Not done |
| 8 | Caffeine 250ppm + EGCG 500 ppm | 5.0 | 70 | 59.0 | Not done |

The percentage reduction in the diameter of the fungal colony in caffeine containing media 2, 4, 5, 6 (500 ppm and 1000 ppm caffeine), 7 and 8 (250 ppm caffeine) was significantly greater when compared to the control medium 1 (Figure 1) and the EGCG containing medium 3 (Table 1). Therefore, the presence of EGCG in the medium apparently does not have an effect on the diameter of the fungal colony while the percentage reduction in the diameter of the fungal colony was less in caffeine media which contained EGCG (media 4, 5 and 6). Therefore, EGCG modulates the inhibitory effect of caffeine on mycelial development on laboratory media.

The presence of EGCG did not significantly modify the effect of caffeine on sporulation. A pronounced reduction in sporulation was observed in the caffeine

containing media when compared to sporulation in the control medium 1 as observed in our previous study⁴ (Table 1). Sporulation in medium 3 (with EGCG) was not significantly different from sporulation in the control medium 1 and the presence of EGCG did not significantly modify the effect of caffeine on sporulation (Table 1, media 7 & 8). A significantly higher number of spores germinated from mycelia grown in EGCG containing media 3 and 4, when compared with caffeine containing media 2, 7 and 8. Therefore, the inhibitory effect of caffeine on spore germination was reduced in the presence of EGCG.

A significant reduction in the weight (56% and 58% respectively) of mycelia was observed in mycelia grown in the EGCG containing media 3 and 4 respectively

(Table 1). A reduction in mycelial growth (23.6%) was also observed in 2, but this difference was not significant when compared with the reduction of mycelial mass observed in the presence of EGCG (3 and 4). The pronounced effect of EGCG on mycelial mass is also clearly seen in Figures 1 and 2, which show that the fungal colony area is more diffuse and less dense in the EGCG containing medium 3 compared to control medium 1, while the fungal colony is more compressed and dense in the caffeine containing media 2 and 5.

Results discussed above were obtained after statistical analysis by one way ANOVA test using SAS computer software. The p value was between 0.004 – 0.0001 for all the experiments. According to the Duncan's Multiple Range test, groups that were significantly different were identified.

Polyphenols such as EGCG are known to have the ability to bind proteins and the restriction in fungal growth may therefore be due to inhibition of certain fungal enzymes⁶. Colonization of xylem vessels by the ambrosia fungus and the presence of globular structures with brown cell contents around the colonized region, reported by us during a previous study⁷ suggest that accumulation of tannins and fungistatic flavanols occurs after gallery formation by SHB attack and may produce an environment that is restrictive to intercellularly spreading pathogens. It is possible that flavanols such as EGCG and other catechins found in tea stems may be involved in a biochemical role to restrict the intercellular spreading of fungal mycelia, and thereby serve to limit the damage caused by SHB to tea stems.

Acknowledgement

The authors thank Prof. N.K.B. Adikaram, Department of Botany, University of Peradeniya, for many useful suggestions; the Tea Research Institute, Hantane sub-station, Kandy for cooperation and assistance and the National Science Foundation of Sri Lanka for research funding.

References

1. Kumar N.S., Hewavitharanage P. & Adikaram N.K.B. (1995). Attack on tea by *Xyleborus fornicatus*: inhibition of the symbiote, *Monacrosporium ambrosium*, by caffeine. *Phytochemistry* **40**: 1113-1116.
2. Hewavitharanage P., Karunaratne S. & Kumar N.S. (1999). Effect of caffeine on shot-hole borer beetle (*Xyleborus fornicatus*) of tea (*Camellia sinensis*). *Phytochemistry* **51**: 35-41.
3. Calnaido D. & Kanapathipillai P. (1967). Tolerance and susceptibility of tea clones to shot hole borer beetle, *Xyleborus fornicatus* Eichh. Coleoptera:Scolytidae. *Tea Science* **38**: 275-281.
4. Karunaratne W.S., Kumar V. & Kumar N.S. (2008). Shot-hole borer infestation of tea: ovicidal effect of caffeine. *Science and Culture* **74**(1-2): 59-64.
5. Arulpragasam P.V. (1992). Disease Control in Asia In: *Tea Cultivation and Consumption* (Eds. K.C. Wilson and M.N. Clifford), pp 555-556, Chapman and Hall, London, UK.
6. Haslam E. (1998). *Practical Polyphenolics. From Structure to Molecular Recognition and Physiological Action*. pp. 323-325 Cambridge University Press, UK.
7. Kumar N.S., Hewavitharanage P. & Adikaram N.K.B. (1998). Histology and fungal flora of shot-hole borer beetle (*Xyleborus fornicatus*) galleries in tea (*Camellia sinensis*). *Journal of the National Science Council of Sri Lanka* **26**(3): 195-207.